

READY FOR ROLVEDON[®]

A Patient and Caregivers Guide
to Education and Home Injection



What is ROLVEDON[®]?

ROLVEDON is a man-made form of granulocyte colony-stimulating factor (G-CSF). G-CSF is a substance produced by the body. It stimulates the growth of neutrophils, a type of white blood cell important in the body's fight against infection. It is not known if ROLVEDON is safe and effective in children.

SELECTED SAFETY INFORMATION

Do not take ROLVEDON if you have had a serious allergic reaction to eflapegrastim, pegfilgrastim or filgrastim products.

ROLVEDON may cause serious side effects, including:

Spleen rupture. Your spleen may become enlarged and can rupture. A ruptured spleen can cause death. Call your healthcare provider right away if you have pain in the left upper stomach-area or your left shoulder.

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).

This is a supplemental guide that supports the training you or a caregiver will have received at your healthcare provider's office.

This guide includes information on the following:

- Why your doctor has prescribed ROLVEDON®
- Understanding neutropenia
- How to store ROLVEDON
- How to inject ROLVEDON
- Common side effects
- Financial support



Not made with natural rubber latex

Always reach out to your healthcare team if you have questions or concerns about injecting ROLVEDON.



Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information, Patient Information, and Instructions for Use.](#)

 **ROLVEDON®**
(eflapegastim-xnst) injection

About ROLVEDON®


ROLVEDON® is an FDA-approved medication used to decrease the incidence of infections associated with chemotherapy.

It is a man-made long-acting form of granulocyte colony-stimulating factor (G-CSF).

What is a G-CSF?

Human G-CSF is a substance your body normally produces that is required to stimulate the production of white blood cells called neutrophils. Neutrophils are essential in fighting off infections. A low neutrophil count increases your risk of serious infections because your body has a reduced ability to fight off harmful microorganisms.

Man-made prescription G-CSFs are similar to human G-CSF as they stimulate the production of neutrophils in the same manner. Man-made G-CSFs can be short-acting or long-acting.



Short-acting man-made GCSFs require multiple injections during a chemotherapy cycle.

Long-acting G-CSFs (LA-GCSFs), such as ROLVEDON, require injection once per chemotherapy cycle.

SELECTED SAFETY INFORMATION

ROLVEDON may cause serious side effects, including:

A serious lung problem called Acute Respiratory Distress Syndrome (ARDS). Call your healthcare provider or get emergency help right away if you have shortness of breath with or without a fever, trouble breathing, or a fast rate of breathing.

Serious allergic reactions. ROLVEDON can cause serious allergic reactions. These reactions can cause a rash all over your whole body, shortness of breath, wheezing, dizziness, swelling around your mouth or eyes, fast heart rate, and sweating. If you have any of these symptoms stop using ROLVEDON and call your healthcare provider or get emergency medical help right away.

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).


 **ROLVEDON®**
(eflapegrastim-xnst) injection

Understanding Neutropenia

Neutropenia is an abnormally low concentration of white blood cells (WBC) in your body. WBCs are a part of your immune system that help prevent and fight off infections, and neutrophils are a type of WBCs that are especially important in this process. Neutropenia occurs when the number of neutrophils in your blood is too low. When this occurs, your body is less able to fight off infections.

Chemotherapy kills cancer cells but can also damage and kill neutrophils leading to neutropenia resulting in an increased risk of infection.

Prescription GCSF medications, like ROLVEDON®, are used to help boost the production of neutrophils and decrease your risk of developing infections associated with chemotherapy.



Chemotherapy

Neutropenia
Low Neutrophil Count

ROLVEDON helps
boost the levels
of neutrophils


Your doctor has prescribed ROLVEDON to help boost the levels of neutrophils in your body to reduce your risk of developing an infection so that you can continue your chemotherapy treatment.

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information, Patient Information, and Instructions for Use.](#)

 **ROLVEDON**[®]
(eflapegrestim-xnst) injection

Febrile Neutropenia

If you develop a fever (greater than 100.4°F or 38.0°C) when your neutrophils are low, you are at an even higher risk of developing a severe infection. This is called **febrile neutropenia**. The consequence of developing febrile neutropenia during chemotherapy treatment varies by person.



How will I know if I have an infection?

- Often the only sign of infection after chemotherapy may be a **fever (100.4 °F [38.0 °C] or higher)**.
- Take your temperature any time you don't feel well or as recommended by your doctor.



If you develop a fever while going through chemotherapy, it is considered a medical emergency, and you must call your healthcare team immediately or go to the emergency room at your local hospital.

SELECTED SAFETY INFORMATION

ROLVEDON may cause serious side effects, including:

Sickle cell crises. You may have a serious sickle cell crisis, which could lead to death, if you have a sickle cell disorder and receive ROLVEDON. Call your healthcare provider right away if you develop symptoms of sickle cell crisis such as pain or difficulty breathing.

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).

 **ROLVEDON**[®]
(eflapegrastim-xnst) injection

Things to Know Before Injecting ROLVEDON®

Before receiving ROLVEDON®, tell your healthcare provider about all of your medical conditions, including if you:

- have a sickle cell disorder
- have kidney problems
- are pregnant or plan to become pregnant. It is not known if ROLVEDON can harm your unborn baby. Tell your healthcare provider right away if you become pregnant during treatment with ROLVEDON.
- are breastfeeding or plan to breastfeed. It is not known if ROLVEDON passes into your breastmilk.



Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.



Do not take ROLVEDON if you have had a serious allergic reaction to eflapegrastim, pegfilgrastim or filgrastim products.

General information about the safe and effective use of ROLVEDON.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ROLVEDON for a condition for which it was not prescribed. Do not give ROLVEDON to other people, even if they have the same symptom that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about ROLVEDON that is written for health professionals.

SELECTED SAFETY INFORMATION

ROLVEDON may cause serious side effects, including:

Kidney injury (glomerulonephritis). ROLVEDON can cause kidney injury. Call your healthcare provider right away if you develop any of the following symptoms:

- swelling of your face or ankles
- blood in your urine or dark colored urine
- you urinate less than usual

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).



ROLVEDON®
(eflapegrastim-xnst) injection


How Will I Receive ROLVEDON®

ROLVEDON® is given as an injection under your skin (subcutaneous injection).

If your healthcare provider decides that the subcutaneous injections can be given at home by you or your caregiver, follow the accompanying “Instructions for Use” on how to prepare and inject a dose of ROLVEDON.

- You and your caregiver will be shown how to prepare and inject ROLVEDON before you use it.
- You will receive 1 injection of ROLVEDON for each cycle of chemotherapy.

Do not attempt to administer ROLVEDON until being properly trained by a healthcare provider.



- You will receive your injection of ROLVEDON about **24 hours after you finish receiving your chemotherapy.**
- You should not receive ROLVEDON for 14 days before or within 24 hours after your dose of chemotherapy.
- If you miss a dose of ROLVEDON, talk to your healthcare provider about when you should receive your next dose.

SELECTED SAFETY INFORMATION

ROLVEDON may cause serious side effects, including:

Increased white blood cell count (leukocytosis). Your healthcare provider will check your blood count during treatment with ROLVEDON.

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).

 **ROLVEDON®**
(eflapegrastim-xnst) injection

How Should I Store ROLVEDON®



Store ROLVEDON® in the refrigerator between 36°F to 46°F (2°C to 8°C) in the **original carton** to protect from light.



Do not freeze ROLVEDON. Throw away (dispose of) any ROLVEDON that has been frozen.



Do not shake ROLVEDON.



Take the carton out of the refrigerator and place the sealed blister tray on a clean flat surface for **at least 30 minutes** to allow it to reach room temperature before use.



Throw away (properly dispose of) any ROLVEDON that has been left at room temperature, 68°F to 77°F (20°C to 25°C), for more than 72 hours.



Keep ROLVEDON and all medicines out of the reach of children.

SELECTED SAFETY INFORMATION

ROLVEDON may cause serious side effects, including:

Decreased platelet count (thrombocytopenia). Your healthcare provider will check your blood during treatment with ROLVEDON. Tell your healthcare provider if you have unusual bleeding or bruising during treatment with ROLVEDON. This could be a sign of decreased platelet counts, which may reduce the ability of your blood to clot.

Capillary Leak Syndrome. ROLVEDON can cause fluid to leak from blood vessels into your body's tissues. This condition is called "Capillary Leak Syndrome" (CLS). CLS can quickly cause you to have symptoms that may become life-threatening. Get emergency help right away if you develop any of the following symptoms:

- swelling or puffiness and are urinating less than usual
- trouble breathing
- swelling of your stomach area (abdomen) and feeling of fullness
- dizziness or feeling faint
- a general feeling of tiredness

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).

 **ROLVEDON®**
(eflapegrastim-xnst) injection


Get to Know the ROLVEDON® Prefilled Syringe



Do not try to give the injection unless you or your caregiver have received training from your healthcare provider.



Make sure to check the name ROLVEDON® appears on the carton and prefilled syringe label and ensure the date on the label has not expired.



Important Information You Need To Know Before Injecting ROLVEDON

- To help reduce needle stick injuries, each prefilled syringe has a needle guard that is automatically activated to cover the needle after the injection is complete.
- **Do not** use the ROLVEDON prefilled syringe if it has been dropped on a hard surface. The prefilled syringe may be broken even if you cannot see the break. Use a new prefilled syringe.
- **Do not** use the ROLVEDON prefilled syringe if the seal on the carton or the seal on the blister tray is broken.
- **Do not** touch the needle guard clips before use. Touching them may cause the needle guard to be activated too early.

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information, Patient Information, and Instructions for Use.](#)

 **ROLVEDON**[®]
(eflapegrastim-xnst) injection

Before Injecting ROLVEDON®

Read the expanded information on each step in the accompanying [Instructions For Use](#) before using ROLVEDON® and each time you get a refill. There may be new information.

This information does not take the place of talking to your healthcare provider about questions regarding your medical condition or treatment.

1

Remove the carton from refrigerator.
Open the carton and remove the blister tray



2

Allow ROLVEDON to reach room temperature



3

Gather the supplies needed for your injection



Alcohol Wipes



Cotton Ball or Gauze Pad



Sharps Container

4

Wash your hands with soap and water



Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).




ROLVEDON®
(eflapeg rastim-xnst) injection

Preparing to Inject ROLVEDON®


5

Remove the prefilled syringe from the blister tray




6

Inspect the prefilled syringe and medicine




7

Select the injection site



8

Clean the injection site




Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).

 **ROLVEDON**[®]
(eflapegrastim-xnst) injection

Ready to Inject ROLVEDON[®]


9

Remove needle cap




10

Insert the needle




11

Inject ROLVEDON[®]



12

Activate the
needle guard



Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).

 **ROLVEDON[®]**
(eflapegrestim-xnst) injection

Disposing of Used Syringes

Put the used prefilled syringe and needle cap in an FDA-cleared sharps disposal container right away after use. Do not throw away (dispose of) the prefilled syringe in your household trash.

If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:

- made of a heavy-duty plastic,
- can be closed with a tight fitting, puncture-resistant lid, without sharps being able to come out,
- upright and stable during use,
- leak-resistant, and
- properly labeled to warn of hazardous waste inside the container.



When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at: <http://www.fda.gov/safesharpsdisposal>.

Do not reuse the prefilled syringes or throw them into household waste.

Do not recycle prefilled syringes or sharps disposal container.

Do not attempt to remove the needle from the prefilled syringe.

Important: Keep the sharps disposal container out of the reach of children.

SELECTED SAFETY INFORMATION

ROLVEDON may cause serious side effects, including:

Myelodysplastic syndrome and acute myeloid leukemia. If you have breast cancer or lung cancer, when ROLVEDON is used with chemotherapy and radiation therapy, or with radiation therapy alone, you may have an increased risk of developing a precancerous blood condition called myelodysplastic syndrome (MDS) or a blood cancer called acute myeloid leukemia (AML). Symptoms of MDS and AML may include tiredness, fever, and easy bruising or bleeding. Call your healthcare provider if you develop these symptoms during treatment with ROLVEDON.

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).

 **ROLVEDON**[®]
(eflapegrastim-xnst) injection

Common Side Effects

The most common side effects of ROLVEDON® include:

- tiredness
- headache
- muscle and joint pain
- nausea
- fever
- back pain
- diarrhea
- anemia
- bone pain
- rash

These are not all the possible side effects of ROLVEDON.

Managing Side Effects: Your healthcare team can help you find ways to manage side effects.

Call your doctor for medical advice about managing side effects.

You may report side effects to FDA at 1-800-FDA-1088.



SELECTED SAFETY INFORMATION

ROLVEDON may cause serious side effects, including:

Inflammation of the aorta (aortitis). Inflammation of the aorta (the large blood vessel which transports blood from the heart to the body) has been reported in patients who received pegfilgrastim products. Symptoms may include fever, abdominal pain, feeling tired, and back pain. Call your healthcare provider if you experience these symptoms.

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).

 **ROLVEDON®**
(eflapegrastim-xnst) injection

Financial Support for ROLVEDON®

ACCESS4ME®

If you need assistance affording your ROLVEDON® medication, we are here to help.

ACCESS4Me® has a dedicated team of reimbursement specialists readily available to assist you and your healthcare provider with the following*:

- Checking your insurance to see if you are covered for ROLVEDON
- Offering a copay program where you may pay as little as \$0 for treatment copays†
- Providing ROLVEDON at no cost for those who may be uninsured or underinsured

* Patients must be enrolled in ACCESS4Me. Patient consent required. Terms and conditions apply. Please contact ACCESS4Me for complete program rules and eligibility requirements.

† Eligibility criteria apply. Patients are not eligible for commercial copay assistance if the prescription is eligible to be reimbursed, in whole or in part, by any state or federal healthcare program.



**For questions about patient support,
please visit ACCESS4Me.com or
call 1-866-582-2737 (866-58-CARES)**

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information, Patient Information, and Instructions for Use.](#)

 **ROLVEDON®**
(eflapegrastim-xnst) injection

Questions About ROLVEDON[®]



Keep this guide and all contents safe so you can refer to them when self-injecting ROLVEDON[®].



Connect with our ACCESS4Me[®] team for financial support.



Call your doctor for questions about this guide and medical advice about managing side effects. You may report side effects to FDA at 1-800-FDA-1088.



Visit MyROLVEDON.com for more information and to view the instructional home injection video

Please see Important Safety Information throughout this brochure and accompanying [full Prescribing Information](#), [Patient Information](#), and [Instructions for Use](#).



ROLVEDON and ACCESS4Me are registered trademarks licensed by Assertio Specialty Pharmaceuticals, LLC. © 2026 Assertio Specialty Pharmaceuticals, LLC. A subsidiary of Assertio Holdings, Inc. All rights reserved.

PP-ROL-00-0872



HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ROLVEDON® safely and effectively. See full prescribing information for ROLVEDON.

ROLVEDON® (eflapegrastim-xnst) injection, for subcutaneous use
Initial U.S. Approval: 2022

RECENT MAJOR CHANGES

Dosage and Administration, Administration (2.2) 07/2025

INDICATIONS AND USAGE

Rolvedon is a leukocyte growth factor indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in adult patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with clinically significant incidence of febrile neutropenia. (1)

Limitations of Use

Rolvedon is not indicated for the mobilization of peripheral blood progenitor cells for hematopoietic stem cell transplantation. (1)

DOSAGE AND ADMINISTRATION

- Recommended Dose: 13.2 mg administered subcutaneously once per chemotherapy cycle. (2.1)
- Administer approximately 24 hours after cytotoxic chemotherapy. Do not administer within the period from 14 days before to 24 hours after administration of cytotoxic chemotherapy. (2.1)

DOSAGE FORMS AND STRENGTHS

Injection: 13.2 mg/0.6 mL solution in a single-dose prefilled syringe. (3)

CONTRAINDICATIONS

Patients with a history of serious allergic reactions to human granulocyte colony-stimulating factors such as eflapegrastim, pegfilgrastim or filgrastim products. (4)

WARNINGS AND PRECAUTIONS

- Fatal splenic rupture: Evaluate patients who report left upper abdominal or shoulder pain for an enlarged spleen or splenic rupture. (5.1)
- Acute respiratory distress syndrome (ARDS): Evaluate patients who develop fever, lung infiltrates, or respiratory distress. Discontinue Rolvedon in patients with ARDS. (5.2)
- Serious allergic reactions, including anaphylaxis: Permanently discontinue Rolvedon in patients with serious allergic reactions. (5.3)
- Sickle Cell Crisis in Patients with Sickle Cell Disorders: Discontinue Rolvedon if sickle cell crisis occurs. (5.4)
- Glomerulonephritis: Evaluate and consider dose-reduction or interruption of Rolvedon if causality is likely. (5.5)
- Leukocytosis: Monitor complete blood count (CBC) during Rolvedon therapy. (5.6)
- Thrombocytopenia: Monitor platelet counts. (5.7)
- Myelodysplastic Syndrome (MDS) and Acute Myeloid Leukemia (AML): Monitor patients with breast and lung cancer using Rolvedon in conjunction with chemotherapy and/or radiotherapy for signs and symptoms of MDS/AML. (5.10)

ADVERSE REACTIONS

The most common adverse reactions (≥20%) are fatigue, nausea, diarrhea, bone pain, headache, pyrexia, anemia, rash, myalgia, arthralgia, and back pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact 1-800-518-1084 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 02/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE	5.12	Nuclear Imaging
2	DOSAGE AND ADMINISTRATION	6	ADVERSE REACTIONS
2.1	Recommended Dosage	6.1	Clinical Trials Experience
2.2	Administration	8	USE IN SPECIFIC POPULATIONS
3	DOSAGE FORMS AND STRENGTHS	8.1	Pregnancy
4	CONTRAINDICATIONS	8.2	Lactation
5	WARNINGS AND PRECAUTIONS	8.4	Pediatric Use
5.1	Splenic Rupture	8.5	Geriatric Use
5.2	Acute Respiratory Distress Syndrome	10	OVERDOSAGE
5.3	Serious Allergic Reactions	11	DESCRIPTION
5.4	Sickle Cell Crisis in Patients with Sickle Cell Disorders	12	CLINICAL PHARMACOLOGY
5.5	Glomerulonephritis	12.1	Mechanism of Action
5.6	Leukocytosis	12.2	Pharmacodynamics
5.7	Thrombocytopenia	12.3	Pharmacokinetics
5.8	Capillary Leak Syndrome	12.6	Immunogenicity
5.9	Potential for Tumor Growth Stimulatory Effects on Malignant Cells	13	NONCLINICAL TOXICOLOGY
5.10	Myelodysplastic Syndrome (MDS) and Acute Myeloid Leukemia (AML) in Patients with Breast and Lung Cancer	13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
5.11	Aortitis	14	CLINICAL STUDIES
		16	HOW SUPPLIED/STORAGE AND HANDLING
		17	PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Rolvedon is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in adult patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with clinically significant incidence of febrile neutropenia.

Limitations of Use

Rolvedon is not indicated for the mobilization of peripheral blood progenitor cells for hematopoietic stem cell transplantation.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

The recommended dosage of Rolvedon is a single subcutaneous injection of 13.2 mg administered once per chemotherapy cycle. Administer approximately 24 hours after cytotoxic chemotherapy. Do not administer within the period from 14 days before to 24 hours after administration of cytotoxic chemotherapy.

2.2 Administration

Rolvedon is administered subcutaneously via a single-dose prefilled syringe.

Prior to use, take the carton out of the refrigerator and place the sealed blister tray on a clean flat surface for a minimum of 30 minutes to allow the product to reach room temperature. Do not warm up the prefilled syringe in any other way. Discard any prefilled syringe left at room temperature for greater than 72 hours. Do not shake. If Rolvedon is accidentally frozen, do not use. Remove the tray from box and carefully remove the prefilled syringe from the tray. If you drop the prefilled syringe onto a hard surface, do not use it. Use a new syringe for the injection.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not administer Rolvedon if discoloration or particulates are observed.

Administer the entire contents of the prefilled syringe.

If the patient or caregiver misses a dose of Rolvedon, instruct them to contact their healthcare provider.

The Rolvedon prefilled syringe does not bear graduation marks and is intended only to deliver the entire contents of the syringe (13.2 mg/0.6 mL) for direct administration.

Not made with natural rubber latex.

3 DOSAGE FORMS AND STRENGTHS

Injection: 13.2 mg/0.6 mL as a clear, colorless, preservative-free solution in a single-dose prefilled syringe.

4 CONTRAINDICATIONS

Rolvedon is contraindicated in patients with a history of serious allergic reactions to eflapegrastim, pegfilgrastim, or filgrastim products. Reactions may include anaphylaxis [*see Warnings and Precautions (5.3)*].

5 WARNINGS AND PRECAUTIONS

5.1 Splenic Rupture

Splenic rupture, including fatal cases, can occur following the administration of recombinant human granulocyte colony-stimulating factor (rhG-CSF) products, such as Rolvedon. Evaluate for an enlarged spleen or splenic rupture in patients who report left upper abdominal or shoulder pain after receiving Rolvedon.

5.2 Acute Respiratory Distress Syndrome

Acute respiratory distress syndrome (ARDS) can occur in patients receiving rhG-CSF products, such as Rolvedon. Evaluate patients who develop fever and lung infiltrates or respiratory distress after receiving Rolvedon for ARDS. Discontinue Rolvedon in patients with ARDS.

5.3 Serious Allergic Reactions

Serious allergic reactions, including anaphylaxis, can occur in patients receiving rhG-CSF products, such as Rolvedon. Permanently discontinue Rolvedon in patients with serious allergic reactions. Rolvedon is contraindicated in patients with a history of serious allergic reactions to eflapegrastim, pegfilgrastim, or filgrastim products [*see Contraindications (4)*].

5.4 Sickle Cell Crisis in Patients with Sickle Cell Disorders

Severe and sometimes fatal sickle cell crises can occur in patients with sickle cell disorders receiving rhG-CSF products, such as Rolvedon. Discontinue Rolvedon if sickle cell crisis occurs.

5.5 Glomerulonephritis

Glomerulonephritis has occurred in patients receiving rhG-CSF products. The diagnoses were based upon azotemia, hematuria (microscopic and macroscopic), proteinuria, and renal biopsy. Generally, events of glomerulonephritis resolved after dose-reduction or discontinuation of rhG-CSF. If glomerulonephritis is suspected, evaluate for cause. If causality is likely, consider dose-reduction or interruption of Rolvedon.

5.6 Leukocytosis

White blood cell (WBC) counts of $100 \times 10^9/L$ or greater have been observed in patients receiving rhG-CSF products. Monitor complete blood count (CBC) during Rolvedon therapy. Discontinue Rolvedon treatment if WBC count of $100 \times 10^9/L$ or greater occurs.

5.7 Thrombocytopenia

Thrombocytopenia has been reported in patients receiving rhG-CSF products. Monitor platelet counts.

5.8 Capillary Leak Syndrome

Capillary leak syndrome has been reported after administration of rhG-CSF products and is characterized by hypotension, hypoalbuminemia, edema and hemoconcentration. Episodes vary in frequency and severity, and may be life-threatening if treatment is delayed. Patients who develop symptoms of capillary leak syndrome should be closely monitored and receive standard symptomatic treatment, which may include a need for intensive care.

5.9 Potential for Tumor Growth Stimulatory Effects on Malignant Cells

The granulocyte colony-stimulating factor (G-CSF) receptor through which Rolvedon acts has been found on tumor cell lines. The possibility that Rolvedon acts as a growth factor for any tumor type, including myeloid malignancies and myelodysplasia, diseases for which Rolvedon is not approved, cannot be excluded.

5.10 Myelodysplastic Syndrome (MDS) and Acute Myeloid Leukemia (AML) in Patients with Breast and Lung Cancer

MDS and AML have been associated with the use of rhG-CSF products in conjunction with chemotherapy and/or radiotherapy in patients with breast and lung cancer. Monitor patients for signs and symptoms of MDS/AML in these settings.

5.11 Aortitis

Aortitis has been reported in patients receiving rhG-CSF products. It may occur as early as the first week after start of therapy. Manifestations may include generalized signs and symptoms such as fever, abdominal pain, malaise, back pain, and increased inflammatory markers (e.g., c-reactive protein and white blood cell count). Consider aortitis in patients who develop these signs and symptoms without known etiology. Discontinue Rolvedon if aortitis is suspected.

5.12 Nuclear Imaging

Increased hematopoietic activity of the bone marrow in response to growth factor therapy has been associated with transient positive bone imaging findings. This should be considered when interpreting bone imaging results.

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are discussed in greater detail in other sections of the labeling:

- Splenic rupture [*see Warnings and Precautions (5.1)*]
- Acute respiratory distress syndrome [*see Warnings and Precautions (5.2)*]
- Serious allergic reactions [*see Warnings and Precautions (5.3)*]
- Sick cell crisis in patients with sickle cell disorders [*see Warnings and Precautions (5.4)*]
- Glomerulonephritis [*see Warnings and Precautions (5.5)*]
- Leukocytosis [*see Warnings and Precautions (5.6)*]
- Thrombocytopenia [*see Warnings and Precautions (5.7)*]
- Capillary leak syndrome [*see Warnings and Precautions (5.8)*]
- Potential for tumor growth stimulatory effects on malignant cells [*see Warnings and Precautions (5.9)*]
- Myelodysplastic Syndrome (MDS) and Acute Myeloid Leukemia (AML) in Patients with Breast and Lung Cancer [*see Warnings and Precautions (5.10)*]
- Aortitis [*see Warnings and Precautions (5.11)*]
- Nuclear Imaging [*see Warnings and Precautions (5.12)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The safety of Rolvedon was evaluated in Study 1 and Study 2 [see *Clinical Studies (14)*]. Patients with early-stage breast cancer received Rolvedon 13.2 mg by subcutaneous injection (n=314) or pegfilgrastim 6 mg by subcutaneous injection (n=326) on Day 2 of each cycle after docetaxel 75 mg/m² and cyclophosphamide 600 mg/m² (TC) chemotherapy.

Among patients receiving Rolvedon, a total of 272 patients received four 21-day treatment cycles. The most common adverse reactions ($\geq 20\%$) were fatigue, nausea, diarrhea, bone pain, headache, pyrexia, anemia, rash, myalgia, arthralgia, and back pain. Table 1 summarizes the adverse reactions that occurred in Studies 1 and 2.

Table 1. Common Adverse Reactions with a Frequency of $\geq 10\%$ Through Week 14 in Patients with Early-Stage Breast Cancer in Study 1 and Study 2

Adverse Reaction	Rolvedon (N = 314) %	Pegfilgrastim** (N=326) %
Fatigue*	181 (58%)	192 (59%)
Nausea	162 (52%)	166 (51%)
Diarrhea	125 (40%)	126 (39%)
Bone pain	119 (38%)	121 (37%)
Headache*	92 (29%)	90 (28%)
Pyrexia*	87 (28%)	84 (26%)
Anemia*	77 (25%)	52 (16%)
Rash*	77 (25%)	99 (30%)
Myalgia	69 (22%)	49 (15%)
Arthralgia	66 (21%)	48 (15%)
Back pain*	63 (20%)	55 (17%)
Decreased appetite	61 (19%)	50 (15%)
Peripheral edema*	57 (18%)	53 (16%)
Abdominal pain*	53 (17%)	67 (21%)
Dizziness*	50 (16%)	38 (12%)
Dyspnea*	49 (16%)	44 (13%)
Cough*	48 (15%)	51 (16%)
Thrombocytopenia*	44 (14%)	17 (5%)
Pain	37 (12%)	42 (13%)
Pain in extremity	36 (11%)	42 (13%)
Local administration reactions*	34 (11%)	27 (8%)
Flushing	32 (10%)	27 (8%)

*Grouped Terms

**Study 1 and Study 2 were not designed to evaluate meaningful comparisons of the incidence of adverse reactions in the Rolvedon and the pegfilgrastim treatment groups.

Permanent discontinuation due to an adverse reaction occurred in 4% of patients who received Rolvedon. The adverse reaction requiring permanent discontinuation in 3 patients who received Rolvedon was rash.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data on Rolvedon use in pregnant women; however, data from published studies with use of other recombinant human granulocyte colony-stimulating factor (rhG-CSF) products in pregnant women have not identified any drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes.

Animal reproduction studies were conducted in rats and rabbits. In rats, eflapegrastim-xnst did not adversely affect embryofetal and/or postnatal development when administered from organogenesis throughout lactation at doses that produced maternal exposures up to 7 times the exposure at the recommended clinical dose. In rabbits, eflapegrastim-xnst caused embryofetal lethality and reduced fetal weight when administered during the organogenesis period at approximately 6 times the exposure at the clinical dose (*see Data*).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risks of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Data

Animal Data

In an embryofetal developmental study in rabbits, eflapegrastim-xnst was administered subcutaneously every other day during the period of organogenesis at doses up to 10 times the clinical exposure at the maximum recommended dose of 13.2 mg. Increased post-implantation loss, reduced number of live fetuses, and reduced fetal body weights were observed at 6 times the clinical exposure, based on AUC. No malformations were observed up to 10 times the clinical exposure, based on AUC.

In an embryofetal developmental study in rats, eflapegrastim-xnst administered subcutaneously every other day during the period of organogenesis did not adversely affect embryofetal development at doses up to 7 times clinical exposure, based on AUC.

In a pre- and post-natal development study in rats, eflapegrastim-xnst administered subcutaneously once weekly from organogenesis through lactation did not adversely affect behavioral, developmental, or reproductive parameters at doses up to 7 times the clinical exposure, based on AUC.

8.2 Lactation

Risk Summary

There are no data on the presence of eflapegrastim-xnst in human milk, the effects on the breastfed child, or the effects on milk production. Endogenous granulocyte colony-stimulating factor (G-CSF) is present in human milk. Other recombinant human granulocyte colony-stimulating factor (rhG-CSF) products are present in human milk at low levels and are not orally absorbed by infants. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Rolvedon and any potential adverse effects on the breastfed child from Rolvedon or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

Of the 314 patients in clinical studies of Rolvedon, 39% were 65 and over, while 6% were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects.

10 OVERDOSAGE

Overdose of Rolvedon may result in leukocytosis and bone pain. In the event of overdose, general supportive measures should be instituted as necessary. Monitor the patient for adverse reactions [see *Adverse Reactions (6)*].

11 DESCRIPTION

Eflapegrastim-xnst is a granulocyte colony-stimulating factor (G-CSF) produced by covalent coupling of a human G-CSF analog (18.6 kDa) and an Fc fragment of human immunoglobulin G4 (IgG4) (49.8 kDa), both derived from recombinant *E. coli*, via a single 3.4 kDa polyethylene glycol linker. The recombinant G-CSF domain in eflapegrastim-xnst is a variant of human G-CSF with two serine substitutions at positions 17 and 65, and no additional N-terminal methionine. Eflapegrastim-xnst has a molecular weight of approximately 72 kDa.

Rolvedon (eflapegrastim-xnst) injection is a sterile, preservative-free, clear, colorless solution supplied in a single-dose prefilled syringe for subcutaneous use. Each 0.6 mL single-dose prefilled syringe contains 13.2 mg of eflapegrastim-xnst, citric acid monohydrate (2.52 mg), mannitol (30 mg), polysorbate 80 (0.72 mg) and sodium chloride (5.26 mg) in Water for Injection. Sodium hydroxide may be used to adjust pH to 5.5 during manufacturing.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Eflapegrastim-xnst is a recombinant human granulocyte growth factor that binds to G-CSF receptors on myeloid progenitor cells and neutrophils, triggering signaling pathways that control cell differentiation, proliferation, migration and survival.

12.2 Pharmacodynamics

Eflapegrastim-xnst has been shown to elevate neutrophil counts in healthy subjects and in cancer patients. Absolute neutrophil count (ANC), C_{max} and area under the effect curve ($AUEC_{last}$) increased with increasing doses of eflapegrastim-xnst in a linear, but less than dose-proportional, manner over a dose range of 45 to 350 mcg/kg.

12.3 Pharmacokinetics

The pharmacokinetics of eflapegrastim-xnst was studied in healthy subjects and patients with breast cancer. After subcutaneous (SC) dosing, the pharmacokinetics of eflapegrastim-xnst was nonlinear and exposure increases were not dose-proportional over the dose range of 45 to 350 mcg/kg.

Absorption

The median T_{max} of eflapegrastim-xnst is 25 hours (6 to 144 hours) in patients with breast cancer following administration of the recommended dosage.

Distribution

The volume of distribution of eflapegrastim-xnst is 1.44 L.

Elimination

The geometric mean half-life of eflapegrastim-xnst in patients with breast cancer is 36.4 hours (Range: 16.1 to 115 hours) during Cycle 1. Eflapegrastim-xnst clearance decreased with increasing doses following single dose administration, suggesting target-mediated clearance of eflapegrastim-xnst by neutrophils. Following repeat administration, clearance increased in Cycle 3 as compared to Cycle 1, potentially due to the subsequent increase in neutrophils. In *in vitro* studies, the IgG4 Fc fragment in eflapegrastim-xnst binds to the neonatal Fc receptor (FcRn), facilitating the FcRn-mediated transcytosis of eflapegrastim-xnst.

Metabolism

Eflapegrastim-xnst is expected to be metabolized by endogenous degradation following receptor-mediated internalization by cells bearing the G-CSF receptor.

Excretion

Eflapegrastim-xnst was not detected in urine.

Drug Interaction Studies

No studies evaluating the drug interaction potential of eflapegrastim-xnst have been conducted.

12.6 Immunogenicity

The observed incidence of anti-drug antibodies is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude meaningful comparisons of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies, including those of eflapegrastim-xnst or of other eflapegrastim products.

Antibodies to eflapegrastim-xnst were detected using bridging enzyme-linked immunosorbent assay (ELISA) with a sensitivity of 65 ng/mL. During the 12-week treatment period in the two randomized studies, twenty-one of 297 (7.1%) patients treated with eflapegrastim-xnst developed antibodies. At the 12-month follow-up visit after the last dose of treatment in the two randomized studies, 28 of 297 (9.4%) patients treated with eflapegrastim-xnst developed antibodies.

Neutralizing antibodies were detected by a cell-based assay with a sensitivity of 1.95 mcg/mL. One patient out of 297 (0.3%) in the eflapegrastim-xnst arm tested positive for neutralizing antibodies post-treatment. Treatment-emergent anti-PEG antibodies were detected by a direct binding ELISA in 126 out of 268 patients (47%) treated with eflapegrastim-xnst. There was no identified clinically significant effect of anti-drug antibodies on pharmacokinetics, pharmacodynamics, safety or effectiveness of Rolvedon over the treatment duration of 12 weeks. There was no identified clinically significant effect of anti-drug antibodies on the safety profile of Rolvedon during the 12-month follow-up period after the last dose.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term carcinogenicity studies have been performed with eflapegrastim-xnst.

Eflapegrastim-xnst was not mutagenic or clastogenic in a standard battery of genotoxicity tests (bacterial mutagenicity (Ames), Chinese hamster ovary cells chromosomal aberration, rat bone marrow micronucleus).

Eflapegrastim-xnst did not affect reproductive performance or fertility in male or female rats at weekly doses up to 7 times the clinical exposure at the maximum recommended dose of 13.2 mg.

14 CLINICAL STUDIES

Patients with Cancer Receiving Myelosuppressive Chemotherapy

The efficacy of Rolvedon to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs was evaluated in two 1:1 randomized, open-label, active-controlled non-inferiority studies of similar design (Study 1 [NCT02643420] and Study 2 [NCT02953340]) that enrolled a total of 643 patients with early-stage breast cancer. Docetaxel 75 mg/m² and cyclophosphamide 600 mg/m² (TC) were administered intravenously every 21 days (on Day 1 of each cycle) for up to 4 cycles. A fixed dose of Rolvedon 13.2 mg/0.6 mL or pegfilgrastim (6 mg/0.6 mL) was administered subcutaneously on Day 2 of each cycle after TC chemotherapy.

The median age of patients enrolled in the two randomized studies was 60 years (Range: 24 to 88), the majority of patients were female (>99%), 77% were White and 12% were Black or African American.

Study 1 enrolled 406 patients; 196 patients to the Rolvedon arm and 210 patients to the pegfilgrastim arm. Study 2 enrolled 237 patients; 118 patients to the Rolvedon arm and 119 patients to the pegfilgrastim arm. Efficacy for both trials was based on the duration of severe neutropenia (DSN) in Cycle 1.

Efficacy results are shown in Table 2. In both studies, Rolvedon was non-inferior to pegfilgrastim. The distributions of the severe neutropenia events in percentage from Cycle 1 for Study 1 and Study 2 are presented in Figure 1.


Table 2. Duration of Severe Neutropenia (DSN) in Cycle 1 (Study 1 and Study 2)

	Study 1		Study 2	
	Rolvedon (n=196)	Pegfilgrastim (n=210)	Rolvedon (n=118)	Pegfilgrastim (n=119)
Mean DSN (SD) (Days)	0.20 (0.503)	0.35 (0.683)	0.31 (0.688)	0.39 (0.949)
Median DSN (Range) (Days)	0 (0, 3)	0 (0, 3)	0 (0, 3)	0 (0, 7)
Difference in DSN (Days)	-0.148		-0.073	
*95% Confidence Interval ^a	-0.265, -0.033		-0.292, 0.129	

^aConfidence intervals were obtained using 2.5 percentile and 97.5 percentile of the 100,000 bootstrap samples with treatment as stratification factor.

*The non-inferiority of Rolvedon to pegfilgrastim was to be declared if the upper bound of 95% CI of the difference in mean DSN between the treatment arms was <0.62 days.

Figure 1. Duration of Severe Neutropenia (DSN) by Day in Cycle 1 (Study 1 and Study 2)



16 HOW SUPPLIED/STORAGE AND HANDLING

Rolvedon (eflapregastim-xnst) injection is a clear, colorless solution supplied in a single-dose prefilled syringe containing 13.2 mg of eflapregastim-xnst in 0.6 mL solution, with 29-gauge 1/2 inch pre-attached (staked) needle with a needle guard.

Rolvedon is provided in a dispensing pack containing one sterile 13.2 mg/0.6 mL prefilled syringe (NDC 82497-132-01).

Store refrigerated at 36°F to 46°F (2°C to 8°C) in the carton to protect from light. Do not shake. Discard any prefilled syringe left at room temperature for greater than 72 hours. Do not freeze; discard syringe if frozen.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information and Instructions for Use). Inform patients to contact their healthcare provider with any questions.

Advise patients of the following risks and potential risks with Rolvedon:

- Splenic rupture and splenomegaly [see *Warnings and Precautions* (5.1)]
- Acute respiratory distress syndrome [see *Warnings and Precautions* (5.2)]
- Serious allergic reactions [see *Warnings and Precautions* (5.3)]
- Sickle cell crisis [see *Warnings and Precautions* (5.4)]
- Glomerulonephritis [see *Warnings and Precautions* (5.5)]
- Leukocytosis [see *Warnings and Precautions* (5.6)]
- Thrombocytopenia [see *Warnings and Precautions* (5.7)]
- Capillary leak syndrome [see *Warnings and Precautions* (5.8)]
- Potential for tumor growth stimulatory effects on malignant cells [see *Warnings and Precautions* (5.9)]
- Increased risk of myelodysplastic syndrome and/or acute myeloid leukemia in patients with breast and lung cancer who receive Rolvedon in conjunction with chemotherapy and/or radiation therapy [see *Warnings and Precautions* (5.10)]
- Aortitis [see *Warnings and Precautions* (5.11)]

Instruct patients who self-administer Rolvedon using the single-dose prefilled syringe of the:

- Importance of following the Instructions for Use
- Dangers of reusing syringes

- Importance of following local requirements for proper disposal of used syringes.

Manufactured by:
Assertio Specialty Pharmaceuticals, LLC
Lake Forest, IL 60045

U.S. License No. 2428

PATIENT INFORMATION
ROLVEDON® (roll veh don)
(eflapegrastim-xnst)
injection

What is Rolvedon?

Rolvedon is a man-made form of granulocyte colony-stimulating factor (G-CSF). G-CSF is a substance produced by the body. It stimulates the growth of neutrophils, a type of white blood cell important in the body's fight against infection. It is not known if Rolvedon is safe and effective in children.

Do not take Rolvedon if you have had a serious allergic reaction to eflapegrastim, pegfilgrastim or filgrastim products.

Before receiving Rolvedon, tell your healthcare provider about all of your medical conditions, including if you:

- have a sickle cell disorder
- have kidney problems
- are pregnant or plan to become pregnant. It is not known if Rolvedon can harm your unborn baby. Tell your healthcare provider right away if you become pregnant during treatment with Rolvedon.
- are breastfeeding or plan to breastfeed. It is not known if Rolvedon passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will I receive Rolvedon?

- Rolvedon is given as an injection under your skin (subcutaneous injection) by a healthcare provider. **If your healthcare provider decides that the subcutaneous injections can be given at home by you or your caregiver, follow the detailed "Instructions for Use" that comes with your Rolvedon for information on how to prepare and inject a dose of Rolvedon.**
- You and your caregiver will be shown how to prepare and inject Rolvedon before you use it.
- You will receive 1 injection of Rolvedon for each cycle of chemotherapy.
- You will receive your injection of Rolvedon about **24 hours after you finish receiving your chemotherapy**.
- You should not receive Rolvedon for 14 days before or within 24 hours after your dose of chemotherapy.
- If you miss a dose of Rolvedon, talk to your healthcare provider about when you should receive your next dose.

What are the possible side effects of Rolvedon?

Rolvedon may cause serious side effects, including:

- **Spleen rupture.** Your spleen may become enlarged and can rupture. A ruptured spleen can cause death. Call your healthcare provider right away if you have pain in the left upper stomach-area or your left shoulder.
- **A serious lung problem called Acute Respiratory Distress Syndrome (ARDS).** Call your healthcare provider or get emergency help right away if you have shortness of breath with or without a fever, trouble breathing, or a fast rate of breathing.
- **Serious allergic reactions.** Rolvedon can cause serious allergic reactions. These reactions can cause a rash all over your whole body, shortness of breath, wheezing, dizziness, swelling around your mouth or eyes, fast heart rate, and sweating. If you have any of these symptoms, stop using Rolvedon and call your healthcare provider or get emergency medical help right away.
- **Sickle cell crises.** You may have a serious sickle cell crisis, which could lead to death, if you have a sickle cell disorder and receive Rolvedon. Call your healthcare provider right away if you develop symptoms of sickle cell crisis such as pain or difficulty breathing.
- **Kidney injury (glomerulonephritis).** Rolvedon can cause kidney injury. Call your healthcare provider right away if you develop any of the following symptoms:
 - swelling of your face or ankles
 - blood in your urine or dark colored urine
 - you urinate less than usual
- **Increased white blood cell count (leukocytosis).** Your healthcare provider will check your blood count during treatment with Rolvedon.
- **Decreased platelet count (thrombocytopenia).** Your healthcare provider will check your blood during treatment with Rolvedon. Tell your healthcare provider if you have unusual bleeding or bruising during treatment with Rolvedon. This could be a sign of decreased platelet counts, which may reduce the ability of your blood to clot.
- **Capillary Leak Syndrome.** Rolvedon can cause fluid to leak from blood vessels into your body's tissues. This condition is called "Capillary Leak Syndrome" (CLS). CLS can quickly cause you to have symptoms that may become life-threatening. Get emergency help right away if you develop any of the following symptoms:
 - swelling or puffiness and are urinating less than usual

- trouble breathing
- swelling of your stomach area (abdomen) and feeling of fullness
- dizziness or feeling faint
- a general feeling of tiredness
- **Myelodysplastic syndrome and acute myeloid leukemia.** If you have breast cancer or lung cancer, when Rolvedon is used with chemotherapy and radiation therapy, or with radiation therapy alone, you may have an increased risk of developing a precancerous blood condition called myelodysplastic syndrome (MDS) or a blood cancer called acute myeloid leukemia (AML). Symptoms of MDS and AML may include tiredness, fever, and easy bruising or bleeding. Call your healthcare provider if you develop these symptoms during treatment with Rolvedon.
- **Inflammation of the aorta (aortitis).** Inflammation of the aorta (the large blood vessel which transports blood from the heart to the body) has been reported in patients who received pegfilgrastim products. Symptoms may include fever, abdominal pain, feeling tired, and back pain. Call your healthcare provider if you experience these symptoms.

The most common side effects of Rolvedon include:

- tiredness
- nausea
- diarrhea
- bone pain
- headache
- fever
- decreased red blood cell count
- rash
- muscle and joint pain
- back pain

These are not all the possible side effects of Rolvedon.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Rolvedon?

- Store Rolvedon in the refrigerator between 36°F to 46°F (2°C to 8°C).
- **Do not** freeze Rolvedon. Throw away (dispose of) any Rolvedon that has been frozen.
- Store Rolvedon in the original carton to protect from light.
- **Do not** shake Rolvedon.
- Take the carton out of the refrigerator and place the sealed blister tray on a clean flat surface for at least 30 minutes to allow it to reach room temperature before use.
- Throw away (dispose of) any prefilled syringe of Rolvedon that has been left at room temperature, 68°F to 77°F (20°C to 25°C), for more than 72 hours.

Keep Rolvedon and all medicines out of the reach of children.

General information about the safe and effective use of Rolvedon.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use Rolvedon for a condition for which it was not prescribed. Do not give Rolvedon to other people, even if they have the same symptom that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about Rolvedon that is written for health professionals.

What are the ingredients in Rolvedon?

Active ingredient: eflapegrastim-xnst

Inactive ingredients: citric acid monohydrate, mannitol, polysorbate 80, and sodium chloride in Water for Injection. Sodium hydroxide may be used to adjust pH to 5.5 during manufacturing.

Manufactured by: Assertio Specialty Pharmaceuticals, LLC, Lake Forest, IL 60045
U.S. License No. 2428


For more information, go to www.rolvedon.com or call 1-800-518-1084.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: 02/2026

INSTRUCTIONS FOR USE
ROLVEDON® [roll veh don]
(eflapegrastim-xnst)
injection, for subcutaneous use
Single-Dose Prefilled Syringe

This Instructions for Use contains information on how to inject Rolvedon. Read this Instructions for Use before using Rolvedon and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment.



Important Information You Need To Know Before Injecting Rolvedon

- It is important that you do not try to give the injection unless you or your caregiver have received training from your healthcare provider.
- Rolvedon is for use under the skin (subcutaneous injection) only.
- To help reduce needle stick injuries, each prefilled syringe has a needle guard that is automatically activated to cover the needle after the injection is complete.
- Make sure the name Rolvedon appears on the carton and prefilled syringe label.
- **Do not** use a Rolvedon prefilled syringe after the expiration date on the label.
- **Do not** use the Rolvedon prefilled syringe if it has been dropped on a hard surface. The prefilled syringe may be broken even if you cannot see the break. Use a new prefilled syringe.
- **Do not** use the Rolvedon prefilled syringe if the seal on the carton or the seal on the blister tray is broken.
- **Do not** shake the Rolvedon prefilled syringe.
- **Do not** touch the needle guard clips before use. Touching them may cause the needle guard to be activated too early.
- **Do not** remove the needle cap until just before you give the injection.

Rolvedon prefilled syringe parts




Figure A

Storing the Rolvedon prefilled syringe

- Store Rolvedon in the refrigerator between 36°F to 46°F (2°C to 8°C).
- **Do not** freeze Rolvedon. Throw away (dispose of) any Rolvedon that has been frozen.
- Store Rolvedon in the original carton to protect from light.
- Take the carton out of the refrigerator and place the sealed blister tray on a clean flat surface for at least 30 minutes to allow it to reach room temperature before use.
- Throw away (dispose of) any prefilled syringe of Rolvedon that has been left at room temperature, 68°F to 77°F (20°C to 25°C), for more than 72 hours.

Keep Rolvedon and all medicines out of the reach of children.

Preparing to inject Rolvedon

Step 1. Remove the carton from refrigerator

- Remove the carton from the refrigerator (see Figure B).
 - Check the expiration date on the carton (see Figure C).
- Do not** use the prefilled syringe if it is expired or if the carton has been previously opened.
- Open the carton and remove the blister tray.




Figure B




Figure C

Step 2. Allow Rolvedon to reach room temperature

Place the sealed blister tray on a clean flat surface and wait 30 minutes to reach room temperature (see Figure D). **Do not** warm up the prefilled syringe in any other way.



Figure D

Step 3. Gather the supplies needed for your injection

Gather the following supplies for the injection (see Figure E):

- 1 sterile cotton ball or gauze pad
- 1 alcohol wipe
- 1 sharps disposal container. See “**Disposing of used Rolvedon prefilled syringes**” at the end of this Instructions for Use.



Figure E

Step 4. Wash your hands

Wash your hands with soap and water.

Step 5. Remove the prefilled syringe from the blister tray

- Open the tray by peeling away the tray cover.
- To remove the prefilled syringe, grip the prefilled syringe by the center of the viewing window as shown in Figure F.
- **Do not** remove the prefilled syringe by the plunger rod, plunger head, or the needle cap.
- **Do not** touch the needle guard clips, as it could activate the needle guard.
- **Do not** remove the needle cap until you are ready to inject.




Figure F

Step 6. Inspect the prefilled syringe and medicine

- Visually inspect the prefilled syringe. **Do not** use the prefilled syringe if it appears damaged (for example, if the syringe is cracked or broken, or if the plunger rod is detached or broken) or if the needle cap is missing.
- Check the expiration date on the prefilled syringe label. **Do not** use if the expiration date has passed.
- Check the medicine inside the prefilled syringe through the viewing window (see Figure G). The medicine inside should be clear and colorless. It is normal to see small air bubbles in the medicine. It does not need to be removed.
- **Do not** use the prefilled syringe if:
 - the medicine appears discolored or cloudy.
 - the medicine contains lumps, flakes, or particles.




Figure G

Step 7. Select the injection site




Figure H

- The recommended injection sites for a subcutaneous injection are the (see Figure H):
 - Abdomen (except for a two-inch area surrounding the navel)
 - Upper outer thigh
 - Outer upper arm (only if someone else is giving you the injection)
 - Upper outer buttock (only if someone else is giving you the injection)
- **Do not** inject into area of the skin that is tender, damaged, bruised, scarred or hard.
- **Do not** inject into moles, scars, birthmarks or stretchmarks.
- If you want to use the same injection site, make sure it is not the same spot on the injection site you used for a previous injection.

Step 8. Clean the injection site

- Clean the injection site with an alcohol wipe (see Figure I).
- Allow the area of skin to dry before injecting.

Do not touch or blow on the cleaned area before injecting.



Figure I

Injecting Rolvedon

Step 9. Remove needle cap

- **Do not** remove the needle cap until you are ready to inject. When ready for the injection, carefully remove the needle cap by pulling it straight off (see Figure J).
- When removing the needle cap, hold the prefilled syringe as shown in Figure J.
- Throw away (dispose of) the needle cap in the sharps disposal container.
- **Do not** put the needle cap back onto the prefilled syringe after it has been removed. This can damage the needle and cause needle stick injuries.
- **Do not** hold the prefilled syringe by the plunger rod or touch the needle guard clips when removing the needle cap.




Figure J

Step 10. Insert the needle

- Hold the prefilled syringe at a 45 to 90-degree angle (see Figure K).
- Using your other hand, gently pinch the cleaned injection site as shown in Figure K.
- Fully insert the needle into the skin in a quick “dart-like” motion.

Do not touch the plunger head while inserting the needle into the skin.




Figure K

Step 11. Inject Rolvedon

- Release the pinched skin.
- Hold the prefilled syringe as shown in Figure L with your thumb on the plunger head ready to inject.
- Slowly press the plunger head down to inject the medicine.
- Hold the syringe steadily in place while the needle is inserted.

Note: You may feel some resistance, this is normal.




Figure L

- Push the plunger head down until it reaches the bottom to make sure all of the medicine is injected (see Figure M).
- You may hear a click when the plunger head reaches the bottom.



Figure M

Step 12. Activate the needle guard

- Gently lift your thumb to release the plunger until the entire needle is covered by the needle guard. Then remove the prefilled syringe from the injection site at the same angle as inserted as shown in Figure N.
- If there is bleeding, lightly press a cotton ball or gauze on the injection site.




Figure N

Disposing of used Rolvedon prefilled syringes

- Put the used prefilled syringe and needle cap in an FDA-cleared sharps disposal container right away after use (see Figure O). Do not throw away (dispose of) the prefilled syringe in your household trash.
- If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:
 - made of a heavy-duty plastic,
 - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
 - upright and stable during use,
 - leak-resistant, and
 - properly labeled to warn of hazardous waste inside the container.
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at: <http://www.fda.gov/safesharpsdisposal>.
- **Do not** reuse the prefilled syringes or throw them into household waste.
- **Do not** recycle prefilled syringes or sharps disposal container.
- **Do not** attempt to remove the needle from the prefilled syringe.

Important: Keep the sharps disposal container out of the reach of children.




Figure O

Manufactured by: Assertio Specialty Pharmaceuticals, LLC Lake Forest, IL 60045

U.S. License No. 2428

For more information, go to www.rolvedon.com or call 1-800-518-1084.

This Instructions for Use has been approved by the U.S. Food and Drug Administration. Issued: 02/2026